



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**198. Proposed by ARTEMAS MARTIN, Washington, D. C.**

Prove that every even number is the sum of two prime numbers.

*Note.*—This problem has long been known and no proof has ever been given. [EDITORS.]

**CORRECTION.**—Number **192**, incorrectly given as **188** in the June issue, page 196, should have the zeros on the right of equations (1), (2), (3) each replaced by  $\square$ , the symbol for a "square number."

## SOLUTIONS OF PROBLEMS.

### ALGEBRA.

**386. Proposed by ELMER SCHUYLER, Brooklyn, N. Y.**

Given the sequence,  $\frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{3}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}, \frac{1}{6}, \frac{5}{6}, \frac{1}{7}, \dots, u_1, u_2, u_3, \dots$ , show that

$$\left[ \frac{\sum_1^n u_i}{n} \right]_{n \rightarrow \infty} = \frac{1}{2}.$$

SOLUTION BY H. A. LEVY, Houghton, Mich.

Consider the sequence  $\frac{1}{q}, \frac{2}{q}, \frac{3}{q}, \dots, \frac{q-1}{q}$ .

(1) Suppose  $q$  to be an odd prime. Then the quantities  $\frac{n}{q}$  ( $n = 1, 2, 3, \dots, q-1$ ) are irreducible fractions, and  $q-1$  must be an even number, say  $2k$ . Then

$$\sum_{n=1}^{q-1} \frac{n}{q} = \frac{1+2+3+\dots+(q-1)}{q} = \frac{q-1}{2} = k.$$

Hence, in such a sequence, there are  $2k$  terms, and the sum of the terms is  $k$ .

(2) Suppose that  $q$  is an even number of the form  $2^r p$ , where  $p$  is any odd prime except one. All the fractions with even numerators are then reducible, and the sequence is

$$\frac{1}{2^r p}, \frac{3}{2^r p}, \frac{5}{2^r p}, \dots, \frac{2^r p - 1}{2^r p}.$$

In this sequence, at least one term is reducible, viz.,  $\frac{p}{2^r p}$ . If there are more,

they are the terms  $\frac{p}{2^r p}, \frac{3p}{2^r p}, \frac{5p}{2^r p}, \dots$  up to the greatest odd multiple of  $p$

less than  $2^r p$ . This odd multiple must therefore be of the form  $4k+1$ . Hence an odd number of terms are reducible. Furthermore, the numerator of the last term of the sequence,  $2^r p - 1$ , is of the form  $4k+1$ , so that there is an odd number of terms in the sequence. Rejecting the reducible terms, therefore, leaves an even number of irreducible terms.

If  $r = 1$ , the number of terms is  $p-1$ , an even number. The sum of the terms is then

$$\left( \frac{1}{2p} + \frac{3}{2p} + \frac{5}{2p} + \cdots + \frac{2p-1}{2p} \right) - \frac{p}{2p} = \frac{p}{2} - \frac{1}{2} = \frac{p-1}{2}.$$

That is, the sum of the terms is half the number of terms. The extension of the proof for the case where  $r > 1$  is obvious.

(3) Suppose  $q$  is the product of two primes, say  $l \cdot m$ . If either  $l$  or  $m$  is 2, this reduces to the case just considered. The sequence is now

$$\frac{1}{lm}, \frac{2}{lm}, \frac{3}{lm}, \cdots, \frac{lm-1}{lm},$$

an even number of terms, since  $lm$  is odd. An even number of terms are reducible. They are

$$\frac{l}{lm}, \frac{2l}{lm}, \frac{3l}{lm}, \cdots, \frac{kl}{lm}, \frac{m}{lm}, \frac{2m}{lm}, \frac{3m}{lm}, \cdots, \frac{k'l}{lm}.$$

$k$  and  $k'$  are both even, because  $k$  is the greatest multiple of  $l$  less than  $m$  and  $k'$  is the greatest multiple of  $m$  less than  $l$ . This leaves an even number of irreducible fractions in the sequence.

The sequence then is

$$\begin{aligned} & \frac{1}{lm} + \frac{2}{lm} + \frac{3}{lm} + \cdots + \frac{lm-1}{lm} \\ & - \left[ \frac{1}{m} + \frac{2}{m} + \frac{3}{m} + \cdots + \frac{m-1}{m} + \frac{1}{l} + \frac{2}{l} + \frac{3}{l} + \cdots + \frac{l-1}{l} \right]. \end{aligned}$$

There are  $lm - l - m + 1$  terms in the sequence, and the sum of the terms is

$$\frac{lm - l - m + 1}{2}.$$

(4) Hence in all cases, the number of irreducible fractions having the same denominator is even, and their sum is half the number of terms.

(5) There remains to be considered one exception, the fractions having 2 as a denominator. Here there is only one term, but the sum of the terms is still one half the number of terms.

(6) Associating together the sequences having 2 as a denominator, 3 as a denominator, 4 as a denominator, etc., we have, besides the term  $\frac{1}{2}$ ,

a sequence of  $2k'$  terms whose sum is  $k'$ ,

a sequence of  $2k''$  terms whose sum is  $k''$ , etc.

Adding these, we have

$$1 + 2(k' + k'' + k''' + \cdots) \text{ terms whose sum is } \frac{1}{2} + k' + k'' + k''' + \cdots.$$

Therefore

$$\frac{\sum_1^m u_i}{n} = \frac{\frac{1}{2} + k' + k'' + k''' + \cdots}{1 + 2(k' + k'' + k''' + \cdots)} = \frac{1}{2}.$$

Also solved by H. C. FEEMSTER.